

**EUR 5194 e**

**COMMISSION OF THE EUROPEAN COMMUNITIES**

**THE HALF-LIFE OF SOME LONG-LIVED ACTINIDES :  
A COMPILATION**

by

**R. VANINBROUKX**

**1974**



**Joint Nuclear Research Centre  
Geel Establishment - Belgium  
Central Bureau for Nuclear Measurements - CBNM**

## LEGAL NOTICE

This document was prepared under the sponsorship of the Commission of the European Communities.

Neither the Commission of the European Communities, its contractors nor any person acting on their behalf:

make any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this document, or that the use of any information, apparatus, method or process disclosed in this document may not infringe privately owned rights; or

assume any liability with respect to the use of, or for damages resulting from the use of any information, apparatus, method or process disclosed in this document.

This report is on sale at the addresses listed on cover page 4

at the price of B.Fr. 50.--
-----------------------------

**Commission of the  
European Communities  
D.G. XIII - C.I.D.  
29, rue Aldringen  
L u x e m b o u r g**

**SEPTEMBER 1974**

This document was reproduced on the basis of the best available copy.

**EUR 5194 e**

THE HALF-LIVE OF SOME LONG-LIVED ACTINIDES : A COMPILATION  
by R. VANINBROUCKX

Commission of the European Communities  
Joint Nuclear Research Centre - Geel Establishment (Belgium)  
Central Bureau for Nuclear Measurements - CBNM  
Luxembourg, September 1974 - 38 Pages - B.Fr. 50,—

A compilation of the reported half-lives of some of the most commonly used long-lived actinides is given. « Best » values and limits of errors are proposed. The compilation shows where further accurate measurements are necessary in order to resolve existing discrepancies

**EUR 5194 e**

THE HALF-LIVE OF SOME LONG-LIVED ACTINIDES : A COMPILATION  
by R. VANINBROUCKX

Commission of the European Communities  
Joint Nuclear Research Centre - Geel Establishment (Belgium)  
Central Bureau for Nuclear Measurements - CBNM  
Luxembourg, September 1974 - 38 Pages - B.Fr. 50,—


A compilation of the reported half-lives of some of the most commonly used long-lived actinides is given. « Best » values and limits of errors are proposed. The compilation shows where further accurate measurements are necessary in order to resolve existing discrepancies

**EUR 5194 e**

THE HALF-LIVE OF SOME LONG-LIVED ACTINIDES : A COMPILATION  
by R. VANINBROUCKX

Commission of the European Communities  
Joint Nuclear Research Centre - Geel Establishment (Belgium)  
Central Bureau for Nuclear Measurements - CBNM  
Luxembourg, September 1974 - 38 Pages - B.Fr. 50,—

A compilation of the reported half-lives of some of the most commonly used long-lived actinides is given. « Best » values and limits of errors are proposed. The compilation shows where further accurate measurements are necessary in order to resolve existing discrepancies



**C O R R I G E N D U M**

**EUR 5194 e      Please replace the title by :**

**"THE HALF-LIFE OF SOME LONG-LIVED ACTINIDES :  
A COMPILATION "**

**on the cover, the bibliographical insert and on the  
title-page.**



**EUR 5194 e**

COMMISSION OF THE EUROPEAN COMMUNITIES

**THE HALF-LIVE OF SOME LONG-LIVED ACTINIDES :  
A COMPILATION**

by

R. VANINBROUKX

1974



**Joint Nuclear Research Centre  
Geel Establishment - Belgium  
Central Bureau for Nuclear Measurements - CBNM**

## **ABSTRACT**

A compilation of the reported half-lives of some of the most commonly used long-lived actinides is given. « Best » values and limits of errors are proposed. The compilation shows where further accurate measurements are necessary in order to resolve existing discrepancies.



## CONTENTS

1. Introduction	5
2. Compilation of reported half-lives	6
2.1 $^{232}\text{U}$	7
2.2 $^{233}\text{U}$	7
2.3 $^{234}\text{U}$	8
2.4 $^{235}\text{U}$	9
2.5 $^{236}\text{U}$	10
2.6 $^{238}\text{U}$	10
2.7 $^{238}\text{Pu}$	10
2.8 $^{239}\text{Pu}$	11
2.9 $^{240}\text{Pu}$	12
2.10 $^{241}\text{Pu}$	12
2.11 $^{242}\text{Pu}$	13
2.12 $^{244}\text{Pu}$	13
2.13 $^{241}\text{Am}$	13
2.14 $^{252}\text{Cf}$	13
3. Conclusions	14
4. References	15
5. Appendix	21



## 1. Introduction

In addition to its general importance an accurate knowledge of the half-lives of the actinides is of particular interest in the quantitative determination, by a radioactivity measurement, of the number of atoms present in a sample (e.g. samples for cross section measurements). Combined with a determination (e.g. by mass spectrometry) or a knowledge of the isotopic composition, the measurement of the radioactivity is the most simple and probably also the most accurate mean for the nondestructive determination of the amount of material in a sample, provided that the decay constants of all the present nuclides are accurately known. Unfortunately, as already stated by Hanna et al. in 1969 <sup>(1)</sup>, several new half-life measurements reported during the last years (e.g. for  $^{233}\text{U}$ ,  $^{234}\text{U}$ ,  $^{241}\text{Pu}$ ,  $^{241}\text{Am}$ ) were in serious disagreement with the previously accepted values. The same authors reported that at that time the half-life of  $^{239}\text{Pu}$  seemed to have escaped so far this malady and they assigned an error of  $\pm 0.2\%$  to the adopted mean value. Two years later a carefully performed new measurement <sup>(2)</sup> yielded a value quoted with an accuracy of  $\pm 0.2\%$  but differing  $1.3\%$  from the previously adopted mean. A further typical example is that of  $^{241}\text{Am}$ . In 1967, as a result of two previously reported values,  $(458.1 \pm 0.5)\text{y}$  <sup>(3)</sup> and  $(457.7 \pm 1.8)\text{y}$  <sup>(4)</sup> a mean of  $458\text{y}$  was adopted <sup>(5)</sup>. In 1967 and 1968 three new values were reported:  $(432.7 \pm 0.6)\text{y}$  <sup>(6)</sup>,  $(436.6 \pm 3.0)\text{y}$  <sup>(7)</sup> and  $(433 \pm 7)\text{y}$  <sup>(8)</sup> and in 1971 a value of  $(433 \pm 2)\text{y}$  was proposed <sup>(9)</sup>. One year later Jove and Robert published a new value of  $(426.3 \pm 2.1)\text{y}$  <sup>(10)</sup>. These examples show that it is very difficult to evaluate reported half-lives. In this report we do not attempt to give an evaluation but rather give a compilation of published data, along with what we consider to be "best" values and limits of errors. One of the

aims of this compilation is to show where high accuracy measurements still seem to be recommended in order to resolve some existing discrepancies, including the apparent systematic differences between values obtained with  $\alpha$ -counting, calorimetric and mass spectrometric techniques.

## 2. Compilation of reported half-lives

Half-life values of some of the most commonly used long-lived actinides (generally only those reported since about 1950) are compiled. The literature data are summarized in the tables A1 to A14 of the appendix. Where available, the material used for the determinations is specified (% of the isotope under investigation relative to the element). The percentages, usually determined by mass spectrometric analysis, are given as quoted by the authors (atomic % or weight %). Furthermore a very brief description of the methods used is given. Unfortunately, often the methods are not well described and important data are missing: e.g. stoichiometry, chemical purity, (both very important when the number of atoms are determined by gravimetric methods), investigation of possible radioactive impurities and their contribution to the counting rates, etc.

The half-life values are given with the errors quoted by the authors. Generally it is difficult, and even impossible, to compare these errors, as often they are not clearly defined. In some cases it is the standard error of the mean of several individual results. In other cases it may be the standard deviation, or the pure statistical error, or a combination of statistical and systematic errors, even summed up in different ways by different authors. It may be  $1\sigma$ ,  $2\sigma$ , or  $3\sigma$ -errors, etc.

## 2.1 $^{232}\text{U}$ (Table A1)

The two reported values, quoted with an error of  $\pm 1.3\%$ , differ by  $2.6\%$ . Since the most recent value is based on a determination of the specific activity of highly enriched material by two independent methods a higher confidence can be attributed to it. Therefore, one can adopt for the half-life of  $^{232}\text{U}$  a value of

$$(72 \pm 2)\text{y}$$

The rather high uncertainty is of less importance for the definition of U-samples by activity measurements. In most U-samples  $^{232}\text{U}$  is only present as a contamination. Due to its high specific activity the contribution of  $^{232}\text{U}$  and daughters to the counting rates can be determined accurately and directly, without a need for an accurate knowledge of the decay constant, by  $\alpha$ - or  $\gamma$ -spectrometric analysis.

## 2.2 $^{233}\text{U}$ (Table A2)

Accurate knowledge of the  $^{233}\text{U}$  half-life is of importance for the determination of fission cross sections. Some earlier experiments will have to be revised when a precise reliable value becomes available <sup>(1)</sup>.  $^{233}\text{U}$  is sometimes useful to spike uranium samples of low specific activity for the purposes of quantitative analysis <sup>(25)</sup>. The  $^{233}\text{U}$ -spike material can be standardized accurately by  $\alpha$ -counting, but an accurate value for the half-life is essential. Unfortunately, the spread of the measured values is many times larger than that expected from the claimed accuracies. Hanna et al. <sup>(1)</sup> estimated from the published data (1952-1969) a best value of  $(1.593 \pm 0.024)10^5\text{y}$ . The reported values can be subdivided into 3 groups.

The first 6 results (1952-1967), all based on  $\alpha$ -counting techniques, yield a mean of  $(1.617 \pm 0.008)10^5$  y. Then (1967-1968) two results yielding a mean of  $(1.554 \pm 0.003)10^5$  y have been published. One of the two values was obtained by  $\alpha$ -counting, the other one by calorimetric techniques. Therefore, the more than 4% lower value can not be explained only by the apparently systematic discrepancy between half-life values obtained by  $\alpha$ -counting and by calorimetry<sup>(2, 19)</sup>. For the third group (1969-1973) three values, all near to  $1.59 \cdot 10^5$  y, are available<sup>(22-24)</sup>. The excellent agreement between these results and especially between the ANL and CBNM values, based on carefully and accurately performed measurements using several independent methods, solves the previously existing discrepancies. From these measurements following "best" value can be adopted:

$$(1.592 \pm 0.003)10^5 \text{ y}$$

### 2.3 $^{234}\text{U}$ (Table A3)

The half-life of  $^{234}\text{U}$  is of great importance since the decay of this isotope is dominant in most uranium samples. Accurate knowledge of the  $^{234}\text{U}$ -decay constant will allow, from activity measurements and knowledge of the isotopic composition, an accurate calculation of the amount of U, especially  $^{235}\text{U}$ , in most of the targets used for cross section measurements. Here also, due to the large spread in reported values, the situation till to 1971 was unsatisfactory. The values reported between 1952 and 1965 yield a mean of  $(2.50 \pm 0.03)10^5$  y. The mean of three new values (1969-1971) is  $(2.443 \pm 0.004)10^5$  y. Since the CBNM<sup>(30)</sup> value is based on a high degree of technique variation (several independent methods, several materials), and is confirmed, within the limits of claimed errors, by the ANL-<sup>(29)</sup> and AECL-<sup>(31)</sup> values one can adopt as "best" value, as already recommended by the 1972 - IAEA Panel on Neutron Standard Reference Data<sup>(87)</sup>

and in agreement with the value selected by De Volpi (20):

$$(2.446 \pm 0.007)10^5 \text{ y}$$

The 0.3% error (3 $\sigma$ -basis) seems, at present, to be satisfactory.

#### 2.4 $^{235}\text{U}$ (Table A4)

For the determination, by activity measurements, of the amount of  $^{235}\text{U}$  in highly enriched U-samples, with very low  $^{234}\text{U}$ -concentrations ( $\leq 0.1\%$ ) the half-life of  $^{235}\text{U}$  is needed. The mean of the values reported before 1971 is  $(7.03 \pm 0.10)10^8 \text{ y}$ .

The stated uncertainty is the standard deviation; it was commonly thought that the accuracy was not better than 2-4% (38, 39). Such a situation is certainly unsatisfactory for accurate cross section determinations.

Fortunately, Jaffey et al. (38) applied improved techniques with the goal of reducing the error to not more than  $\pm 0.2\%$ . The result of their carefully performed measurements was  $(7.038 \pm 0.005)10^8 \text{ y}$ , thus agreeing within 0.1% with the mean mentioned above. The quoted error (less than 0.1%) however is only the statistical standard error of the mean based on the scatter of the observed data. Employing the same error philosophy as for the CBNM-measurements on  $^{234}\text{U}$  (30) by adding to the 3 $\sigma$ -statistical error an upper limit of  $0.005 \cdot 10^8 \text{ y}$  for possible systematic effects, as estimated by the authors, one gets an error of  $\pm 0.3\%$ . Since this result is obtained from the most accurate measurements one should (at least as long as no further accurate measurements are reported) adopt as "best" value

$$(7.038 \pm 0.020)10^8 \text{ y}$$

In order to be sure that possible systematic effects are not

underestimated a second high accuracy determination, preferably with several independent methods, is desirable.

## 2.5 $^{236}\text{U}$ (Table A5)

The half-life of  $^{236}\text{U}$  is of less importance for the definition by activity measurements of commonly used U-samples, where only small concentrations of  $^{236}\text{U}$  are present.

The reported values are spread over about 5%. According to Flynn et al. <sup>(41)</sup> likely sources of systematic errors could explain the deviation of the first result in Table A5 but no explanation could be given for the discrepancy between the two other values. Therefore it is not very meaningful to propose a best value as long as no new accurate measurements are performed. As long as results of such measurements are not available one can adopt the most accurate result, but with an appreciably higher uncertainty:

$$(2.34 \pm 0.02)10^7 \text{ y}$$

## 2.6 $^{238}\text{U}$ (Table A6)

Jaffey et al. <sup>(38)</sup> performed careful measurements on highly depleted material. The  $^{238}\text{U}$ -activity of the two samples used was respectively 90.7% and 99.7% of the total  $\alpha$ -activity. Since the obtained value for the half-life is confirmed by another precise measurement <sup>(45)</sup> one can adopt the ANL-value as "best" value, but with an increased error, as obtained for  $^{235}\text{U}$  (3 $\sigma$  statistical+ systematic error):

$$(4.468 \pm 0.010)10^9 \text{ y}$$

## 2.7 $^{238}\text{Pu}$ (Table A7)

Little information is available. Because of the uncertainties on the Cm-isotopic composition and the  $^{242}\text{Cm}$  half-life for the



method based on the growth of  $^{238}\text{Pu}$  (46) one should attribute a higher confidence to the results obtained by the calorimetric method, however with a considerably higher uncertainty as quoted, mainly because of the generally existing discrepancy between results obtained by calorimetric methods and counting techniques. Several figures between 87.4y and 87.8y, all attributed to calorimetric determinations performed at Mound Laboratory, are reported without details (47-50). It seems likely that the more recent values are revised figures of the previous ones. As long as no new accurate measurements are performed one can adopt for the half-life of  $^{238}\text{Pu}$ :

$$(87.8 \pm 0.8)\text{y}$$

## 2.8 $^{239}\text{Pu}$ (Table A8)

The half-life of  $^{239}\text{Pu}$  is very important for fission cross section determinations. Before the 1971-value of Oetting (2) was published the situation seemed to be very satisfactory. The mean of 8 values (19, 52-58) (including one calorimetric determination (19)) was  $(2.4365 \pm 0.0041)10^4\text{y}$ . The mean of 4 values (19, 56-58) quoted by the authors with an accuracy of  $\pm 0.2\%$  or better was also  $(2.4365 \pm 0.0044)10^4\text{y}$ . Only one calorimetric value (51), with a claimed error of 1%, deviated 1.1% from the mean. The situation was completely perturbed by the recent calorimetric determination of Oetting on a very pure material, yielding a half-life value (claimed to be 0.2% accurate) which is 1.3% lower than the mentioned mean. High accuracy measurements are urgently needed to resolve this discrepancy. Measurements will be started soon at CBNM. Until new data are available one can adopt as half-life:

$$(2.430 \pm 0.025)10^4\text{y}$$

## 2.9 $^{240}\text{Pu}$ (Table A9)

The reported values, with a mean of  $6.52 \cdot 10^3 \text{ y}$ , are spread over 8%, although most values are claimed to be better than 1% accurate.

The mean of 4 values, (19, 54, 57, 61) quoted with an error less than 1%, is  $(6.62 \pm 0.10)10^3 \text{ y}$ . Combining both means with the precise calorimetric value of Oetting one can adopt:

$$(6.55 \pm 0.07)10^3 \text{ y}$$

Accurate measurements should be performed in order to reduce the uncertainty.

## 2.10 $^{241}\text{Pu}$ (Table A10)

Here we have a very particular situation. Four measurements (64-67), using the  $^{241}\text{Am}$ -growth method, yield a mean of  $(14.0 \pm 0.15) \text{ y}$ . Three mass spectrometric measurements (69-73) yield  $(14.9 \pm 0.25)$ , the mean of the two most precise measurements (70-73) being  $(15.0 \pm 0.1) \text{ y}$ . One measurement (69) based on the changes of the reactivity yields a value between both means.

The discrepancies of up to 7% between the different methods, especially the growth- and mass spectrometric method, should be resolved by careful measurements using all available independent methods. As long as results of such measurements are not available one should adopt:

$$(14.5 \pm 0.5) \text{ y}$$

The 0.34y isomeric state reported by Stepan and Nisle (68, 74) was not observed by other investigators (70, 75).

2.11  $^{242}\text{Pu}$  (Table A11)

The spread of the reported values, with a mean of  $3.77 \cdot 10^5 \text{ y}$  is 6%. The mean of the two values determined relative to the half-lives of  $^{240}\text{Pu}$  and  $^{239}\text{Pu}$  is  $(3.87 \pm 0.01)10^5 \text{ y}$ . Since the error on the reference half-lives is not considerably higher than 1% one can adopt for the  $^{242}\text{Pu}$ -half-life:

$$(3.87 \pm 0.05)10^5 \text{ y}$$

2.12  $^{244}\text{Pu}$  (Table A12)

The mean of the two values measured on enriched material, relative to  $^{239}\text{Pu}$ ,  $^{240}\text{Pu}$ , and  $^{242}\text{Pu}$ , is  $(8.23 \pm 0.05)10^7 \text{ y}$ . On the basis of an uncertainty of about 1% on the reference half-lives one can adopt

$$(8.2 \pm 0.1)10^7 \text{ y}$$

2.13  $^{241}\text{Am}$  (Table A13)

Comments have been given in the introduction. One can adopt the mean of the four recent determinations:

$$(432 \pm 4) \text{ y}$$

2.14  $^{252}\text{Cf}$  (Table A14)

Since 1965 four accurate values (quoted errors of 0.2 to 0.4%) have been published for the half-life of  $^{252}\text{Cf}$ , a nuclide often used as a standard for neutron sources. One can adopt the mean of these values:

$$(2.64 \pm 0.02) \text{ y}$$

### 3. Conclusions

From the available data "best" values are proposed, together with estimated uncertainties. These figures are summarized in table 1. If accuracies of 0.2-0.3% would be pursued a lot of new accurate determinations should be performed. The most urgent measurements are the Pu-isotopes (for which the half-lives are not better known than to about 1%), especially  $^{239}\text{Pu}$  and  $^{241}\text{Pu}$ , and  $^{252}\text{Cf}$ . Less urgent, but still interesting, are  $^{232}\text{U}$ ,  $^{235}\text{U}$ ,  $^{236}\text{U}$ , and  $^{241}\text{Am}$ . Special efforts should be made in order to resolve the discrepancies between the values obtained with  $\alpha$ -counting, calorimetric and mass spectrometric techniques.

Table 1: Proposed half-life values

Nuclide	Half-life
$^{232}\text{U}$	$(72 \pm 2)\text{y}$
$^{233}\text{U}$	$(1.592 \pm 0.003)10^5\text{y}$
$^{234}\text{U}$	$(2.446 \pm 0.007)10^5\text{y}$
$^{235}\text{U}$	$(7.038 \pm 0.020)10^8\text{y}$
$^{236}\text{U}$	$(2.34 \pm 0.02)10^7\text{y}$
$^{238}\text{U}$	$(4.468 \pm 0.010)10^9\text{y}$
$^{238}\text{Pu}$	$(87.8 \pm 0.8)\text{y}$
$^{239}\text{Pu}$	$(2.430 \pm 0.025)10^4\text{y}$
$^{240}\text{Pu}$	$(6.55 \pm 0.07)10^3\text{y}$
$^{241}\text{Pu}$	$(14.5 \pm 0.5)\text{y}$
$^{242}\text{Pu}$	$(3.87 \pm 0.05)10^5\text{y}$
$^{244}\text{Pu}$	$(8.2 \pm 0.1)10^7\text{y}$
$^{241}\text{Am}$	$(432 \pm 4)\text{y}$
$^{252}\text{Cf}$	$(2.64 \pm 0.02)\text{y}$

#### 4. References

1. G.C. Hanna, C.H. Westcott, H.D. Lemmel, B.R. Leonard, Jr., J.S. Story, and P.M. Attree, Atomic Energy Rev. 7, 3 (1969)
2. F.L. Oetting, R.F.P-1469 (1971)
3. G.R. Hall and T.L. Markin, J. Inorg. Nucl. Chem. 4, 137 (1957)
4. J.C. Wallmann, P. Graf, and L. Goda, J. Inorg. Nucl. Chem. 7, 199 (1958)
5. C.M. Lederer, J.M. Hollander, and I. Perlman, Table of Isotopes, 6th Ed., John Wiley and Sons, New York (1967)
6. F.L. Oetting and S.R. Gunn, J. Inorg. Nucl. Chem. 29, 2659 (1967)
7. R.E. Stone and E.K. Hulet, J. Inorg. Nucl. Chem. 30, 2003 (1968)
8. L.C. Brown and R.C. Probst, J. Inorg. Nucl. Chem. 30, 2591 (1968)
9. Y.A. Ellis, Nuclear Data Sheets B6 , 621 (1971)
10. J. Jove and R. Robert, Radiochem. Radioanal. Lett. 10, 139 (1972)
11. P.A. Sellers, C.M. Stevens, and M.H. Studier, Phys. Rev. 94, 952 (1954)
12. J.M. Chilton, R.A. Gilbert, R.E. Leuze, and W.S. Lyon, J. Inorg. Nucl. Chem. 26, 395 (1964)
13. E.K. Hyde, in "Production and separation of  $^{233}\text{U}$ ", p. 16, Ed. L.I. Katzin, TID-5223 (1952)
14. P.A. Sellers, W.C. Bentley, and M.H. Studier, ANL-5411, 10 (1955)

15. C.B. Bigham, G.C. Hanna, P.R. Tunnicliffe, P.J. Campion, M. Lounsbury, and D.R. MacKenzie, Second Int. Conf. Peaceful Uses Atom. Energy (Proc. Conf. Geneva 1958) 16, 125, U.N. New York (1959)
16. Y.P. Dokuchaev and I.S. Osipov, Atomnaya Energiya 6, 73 (1959)
17. D.S. Popplewell, J. Nucl. Energy 14, 50 (1961)
18. H.R. Ihle, E. Langenscheidt, and A.P. Murrenhoff, Jul-491-PC (1967)
19. F.L. Oetting, in "Thermodynamics of Nuclear Materials, 1967" IAEA, Vienna (1968)
20. A. De Volpi, ANL-7830 (1971)
21. R.L.G. Keith, J. Nucl. Energy 22, 471 (1968)
22. R.W. Durham, AECL Progress Report, Chemistry and Materials Division, Sept. (1969)
23. A.H. Jaffey, K.F. Flynn, W.C. Bentley, and J.O. Karttunen, Phys. Rev. C9, 1991 (1974)
24. R. Vaninbroukx, et al. to be published (1974)
25. R.L.G. Keith, A. McNair, and A.L. Rodgers, J. Nucl. Energy 22, 477 (1968)
26. C.A. Kienberger, Phys. Rev. 87, 520 (1952)
27. E.H. Fleming, Jr, A. Ghiorso, and B.B. Cunningham, Phys. Rev. 88, 642 (1952)
28. P.H. White, G.J. Wall, and F.R. Pontet, J. Nucl. Energy 19, 33 (1965)
29. J.W. Meadows, ANL-7610, p. 44 (1969)

30. P. De Bièvre, K.F. Lauer, Y. Le Duigou, H. Moret, G. Müschenborn, J. Spaepen, A. Spornol, R. Vaninbrouckx, and V. Verdingh, Intern. Conf. on Chemical Nuclear Data (Proc. Conf. Canterbury 1971), p. 221, Institution of Civil Engineers, London (1971)
31. M. Lounsbury and R.W. Durham, Intern. Conf. on Chemical Nuclear Data (Proc. Conf. Canterbury 1971) p. 215, Institution of Civil Engineers, London (1971)
32. A.O. Nier, Phys. Rev. 55, 150 (1939)
33. G.B. Knight, K-663, ORNL (1950)
34. G.J. Sayag, Comptes Rendus 232, 2091 (1951)
35. E. Würger, K.P. Meyer, and P. Huber, Helv. Phys. Acta 30, 157 (1957)
36. A.J. Deruytter, I.G. Schröder, and J.A. Moore, Nucl. Sci. Eng. 21, 325 (1965)
37. P.O. Banks and L.T. Silver, J. Geophys. Res. 71, 4037 (1966)
38. A.H. Jaffey, K.F. Flynn, L.E. Glendenin, W.C. Bentley, and A.M. Essling, Phys. Rev. C4, 1889 (1971)
39. A. Artna-Cohen, Nuclear Data Sheets B6, 287 (1971)
40. A.H. Jaffey, H. Diamond, A. Hirsch, and J. Mech, Phys. Rev. 84, 785 (1951)
41. K.F. Flynn, A.H. Jaffey, W.C. Bentley, and A.M. Essling, J. Inorg. Nucl. Chem. 34, 1121 (1972)
42. C.A. Kienberger, Phys. Rev. 76, 1561 (1949)
43. A.F. Kovarik and N.I. Adams, Jr., Phys. Rev. 98, 46 (1955)
44. R.B. Leachman and H.W. Schmitt, J. Nucl. Energy 4, 38 (1957)

45. J. Steyn and F.W. Strelow, in "Metrology of Radionuclides" p. 155, IAEA, Vienna (1960)
46. D.C. Hoffman, G.P. Ford, and F.O. Lawrence, J. Inorg. Nucl. Chem. 5, 6 (1957)
47. J.F. Eichelberger, G.R. Grove, and L.V. Jones, MLM-1238 (1965)
48. K.C. Jordan, MLM-1443 (1967)
49. R. Sher, BNL-50233 (1970)
50. R.A. Benson (1969), reported by Y.A. Ellis in Nuclear Data Sheets, B4, 635 (1970)
51. J.W. Stout and W.M. Jones, Phys. Rev. 71, 582 (1947)
52. B.B. Cunningham and L.B. Werner, J. Am. Chem. Soc. 71, 1521 (1949)
53. E.F. Westrum, Jr, J.C. Hindman, and R. Greenlee, in "The Transuranium Elements" Ed. G.T. Seaborg, J. Katz, and W.M. Manning, McGraw-Hill, New York (1949)
54. J.C. Wallmann, UCRL-1255 (1951)
55. G.W. Farwell, J.E. Roberts, and A.C. Wahl, Phys. Rev. 94, 363 (1954)
56. T.L. Markin, J. Inorg. Nucl. Chem. 9, 320 (1959)
57. Y.P. Dokuchaev, Atomnaya Energiya 6, 74 (1959)
58. H.R. Ihle, A.P. Murrenhoff, and M. Karayannis, Jül-347-PC (1966) and in "Standardization of Radionuclides", p. 69, IAEA, Vienna (1967)
59. F.L. Oetting, J. Inorg. Nucl. Chem. 27, 2151 (1965)
60. E.F. Westrum, Jr, Phys. Rev. 83, 1249 (1951)
61. M.G. Inghram, D.C. Hess, P.R. Fields, and G.L. Pyle, Phys. Rev. 83, 1250 (1951)



62. J.P. Butler, T.A. Eastwood, T.L. Collins, M.E. Jones, F.M. Rourke, and R.P. Schuman, Phys. Rev. 103, 634 (1956)
63. M.R. Schmorak, Nuclear Data Sheets B6, 661 (1971)
64. D.R. MacKenzie, M. Lounsbury, and A.W. Boyd, Phys. Rev. 90, 327 (1953)
65. B. Rose and J. Milsted, J. Nucl. Energy 2, 264 (1956)
66. F. Brown, G.G. George, D.E. Green, and D.E. Watt, J. Inorg. Nucl. Chem. 13, 192 (1960)
67. H.L. Smith, J. Inorg. Nucl. Chem. 17, 178 (1961)
68. R.G. Nisle and I.E. Stepan, Nucl. Sci. Eng. 39, 257 (1970)
69. W.R. Shields, N.B.S. Techn. Note 546, p. 25 (1970)
70. M.J. Cabell and M. Wilkins, J. Inorg. Nucl. Chem. 33, 903 (1971)
71. M.J. Cabell, Intern. Conf. on Chemical Nuclear Data (Proc. Conf. Canterbury 1971), p. 189, Institution of Civil Engineers, London (1971)
72. M.J. Cabell, J. Inorg. Nucl. Chem. 30, 2583 (1968)
73. R.K. Zeigler and Y. Ferris, J. Inorg. Nucl. Chem. 35, 3417 (1973)
74. I.E. Stepan and R.G. Nisle, Trans Am. Nucl. Soc. 9, 451 (1966)
75. C.E. Bemis, Jr, R.J. Silva, J.E. Bigalow, and A.M. Friedman, Inorg. Nucl. Chem. Lett. 6, 747 (1970)
76. J.P. Butler, M. Lounsbury, and J. Merritt, Can. J. Chem. 34, 253 (1956)
77. J.F. Mech, H. Diamond, M.H. Studier, P.R. Fields, A. Hirsch, C.M. Stevens, R.F. Barnes, D.J. Henderson, J.R. Huizenga, Phys. Rev. 103, 340 (1956)

78. C.E. Bemis, Jr, J. Halperin, and R. Eby, J. Inorg. Nucl. Chem. 31, 599 (1969)
79. R.W. Durham and F. Malson, Can. J. Phys. 48, 716 (1970)
80. H. Diamond and R.F. Barnes, Phys. Rev. 101, 1064 (1956)
81. P.R. Fields, A.M. Friedman, J. Milsted, J. Lerner, C.M. Stevens, D. Metta, and W.K. Sabine, Nature 212, 131 (1966)
82. T.A. Eastwood, J.P. Butler, M.J. Cabell, H.G. Jackson, R.P. Schuman, F.M. Rourke, and T.L. Collins, Phys. Rev. 107, 1635 (1957)
83. D. Metta, H. Diamond, R.F. Barnes, J. Milsted, J. Gray, Jr, D.J. Henderson, and C.M. Stevens, J. Inorg. Nucl. Chem. 27, 33 (1965)
84. A. De Volpi and K.G. Porges, Inorg. Nucl. Chem. Lett. 5, 699 (1969)
85. B.J. Mijnheer and E. Van den Hauten-Zuidema, Intern. J. Appl. Rad. Isotopes 24, 185 (1973)
86. V. Spiegel, Nucl. Sci. Engin. 53, 326 (1974)
87. I.A.E.A., Summaries, Conclusions and Recommendations from the 2nd Panel on Neutron Standard Reference Data, 20-24 Nov. (1972)

5. Appendix

Tables A1 to A14 summarize the most interesting literature data.

Material : % isotope investigated means the abundance, in atomic % or weight % (as given by the authors), of the isotope for which the half-life is determined.

Method : U-determination by isotopic dilution means always mass spectrometric isotopic dilution analysis.

Table A1. Half-life of  $^{232}\text{U}$

Author (Year)	Material % isotope investig.	Method	Half-life in units of one y	Remarks
Sellers et al. (1954) <sup>(11)</sup>	0.14	$2\pi\alpha$ -counting; $^{232}\text{U}$ determination by isotopic dilution mass spectro- metric analysis	$73.6 \pm 1.0$	
Chilton et al. (1964) <sup>(12)</sup>	99.2	calorimetry and $2\pi\alpha$ -counting; U-determ. by coulometry	$71.7 \pm 0.9$	

Table A2. Half-life of  $^{233}\text{U}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^5$ y	Remarks
Hyde (1952) <sup>(13)</sup>	96.4	2 $\pi\alpha$ -counting; gravimetric U-determ.	$1.62 \pm 0.01$	recalculated by Hanna <sup>(1)</sup> from the ratio $\lambda_{\text{nat U}}/\lambda_{^{233}\text{U}}$
Sellers et al. (1955) <sup>(14)</sup>		$\alpha$ -activity determin. with counters of known geometry; gravimetric U-determin.	$1.615 \pm 0.004$	
Bigham et al. (1959) <sup>(15)</sup>	99.7	2 $\pi\alpha$ -counting; U-determin. by isotopic dilution; radiochemical purity analysis by $\alpha$ -spectrometry	$1.603 \pm 0.008$	
Dokuchaev and Osipov (1959) <sup>(16)</sup>	~ 95	low geom. $\alpha$ -counting; gravimetric U-determ.; radiochem. purity analysis by $\alpha$ -spectrometry	$1.626 \pm 0.008$	
Popplewell (1961) <sup>(17)</sup>	99.8	low geom. $\alpha$ -counting; gravimetric U-determ.	$1.615 \pm 0.009$	original value of 1.540 corrected for $^{232}\text{U}$ daugh- ters as quoted by De Volpi <sup>(20)</sup>
Ihle et al. (1967) <sup>(18)</sup>	87.8	liquid scintill. $\alpha$ -counting; gravim. U-determ.; radiochem. purity analysis by $\alpha$ spectrometry	$1.621 \pm 0.003$	
Oetting (1968) <sup>(19)</sup>	96.9	calorimetry; gravimetric U determ.	$1.554 \pm 0.003$	
Keith (1968) <sup>(21)</sup>	> 99.9	low geometry $\alpha$ -counting; U-determ. by isotopic dilution	$1.553 \pm 0.010$	
continued on next page				

Table A2-bis.

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^5$ y	Remarks
Durham (1969) (22)	99.5	low geom. $\alpha$ -counting	$1.583 \pm 0.007$	the quoted error is only the standard error
Jaffey et al. (1974) (23)		intermed. geom. $\alpha$ -counting; U- determ. by titration; radiochem. purity analysis by $\alpha$ -spectrometry	$1.5911 \pm 0.0015$	
Vaninbrouckx et al. (1974) (24)	> 99.9	$4\pi\alpha$ -, liq. scint. and low geom. - $\alpha$ counting; U-determ. by coulometry and isotopic dilution; radiochem. purity analysis by $\alpha$ - and $\gamma$ -spectro- metry	$1.5925 \pm 0.0040$	

Table A3. Half-life of  $^{234}\text{U}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^5 \text{ y}$	Remarks
Kienberger (1952) <sup>(26)</sup>	94.7	$2\pi\alpha$ -counting; gravimetric U-determ.; chemical purity determ. by spectrographic analysis	$2.520 \pm 0.008$	according to De Volpi (20) error should be $0.033 \cdot 10^5 \text{ y}$
Fleming et al. (1952) <sup>(27)</sup>	96.0	intermed. geom. $\alpha$ -counting; gravimetric U-determ.; chemical purity by spectrographic analysis	$2.475 \pm 0.016$	
Bigham et al. (1959) <sup>(15)</sup>	0.6	$2\pi\alpha$ -counting; U-determ. by isotopic dilution; radiochem. analysis by $\alpha$ spectrometry	$2.519 \pm 0.025$	calculated by Hanna (1) from data on "enriched U"
White et al. (1965) <sup>(28)</sup>	1.2	low geom. $\alpha$ -counting; U-determ. by gravimetry, spectrophotometry, and coulometry; radiochemical analysis by $\alpha$ -spectrometry	$2.47 \pm 0.03$	
Meadows (1969) <sup>(29)</sup>	0.7-1.1	$2\pi\alpha$ - and low geom. $\alpha$ -counting; U-determ. by titration	$2.439 \pm 0.014$	
De Bièvre et al. (1971) <sup>(30)</sup>	0.2-99.9	$4\pi\alpha$ -, liq. scint., and low geom. $\alpha$ -counting; U-determ. by gravimetry, coulometry and isotopic dilution; radiochemical analysis by $\alpha$ -spectrometry	$2.446 \pm 0.007$	
Lounsbury and Durham (1971) <sup>(31)</sup>	1.1	low geom. $\alpha$ -counting; U-determ. by isotopic dilution	$2.444 \pm 0.012$	

Table A4. Half-life of  $^{235}\text{U}$ 

Author (Year)	Material % isotope investig.	Method $R = \frac{\text{activity } ^{235}\text{U}}{\text{activity } ^{238}\text{U}}$ $R' = \frac{\text{activity } ^{235}\text{U}}{\text{activity } ^{234}\text{U}}$	Half-life in units of $10^8 \text{ y}$	Remarks
Nier (1939) (32)	nat U	measurement of R from Pb/U-ratios	7.04 $\pm$ 0.15	recalculated value, cited by Jaffey (36)
Knight (1950) (33)	"enriched $^{235}\text{U}$ "	measurement of the specific activity	7.10 $\pm$ 0.16	"
Sayag (1951) (34)	nat U	measurement of R and R' by ionization chamber energy analysis	6.94 $\pm$ 0.25	"
Fleming et al. (1952) (27)	99.9	intermed. geom. $\alpha$ -counting; gravimetric U-determ.; chem. purity by spectrogr. analysis	7.13 $\pm$ 0.16	
Würger et al. (1957) (35)	nat U	measurement of R and R' by ion chamber and $\alpha$ - $\gamma$ coincidence counting	6.84 $\pm$ 0.15	recalculation by Jaffey (36) yields $6.92 \cdot 10^8 \text{ y}$
White et al. (1965) (28)	99.3	low geom. $\alpha$ -counting; U-determ. by gravimetry, spectrophotometry, and coulometry; radiochem. analysis by $\alpha$ -spectrometry	7.13 $\pm$ 0.09	
Deruytter et al. (1965) (36)	nat U	measurement of R and R' using solid state detectors	6.92 $\pm$ 0.09	recalculation by Jaffey (38) yields $6.97 \cdot 10^8 \text{ y}$
continued on next page				



Table A4-bis.

Author (Year)	Material % isotope investig.	Method $R = \frac{\text{activity } ^{235}\text{U}}{\text{activity } ^{238}\text{U}}$ $R' = \frac{\text{activity } ^{235}\text{U}}{\text{activity } ^{234}\text{U}}$	Half-life in units of $10^8$ y	Remarks
Banks and Silver (1966) <sup>(37)</sup>	nat U	measurement of R by mass spectro- metric determination of $^{207}\text{Pb}/^{206}\text{Pb}$	$7.09 \pm ?$	recalculation by Jaffey <sup>(38)</sup> yields $7.02 \cdot 10^8$ y
Jaffey et al. (1971) <sup>(38)</sup>	> 99.9	intermed. geom. $\alpha$ -counting; U-de- term. by titration; radiochem. purity by $\alpha$ -spectrometry	$7.038 \pm 0.005$	quoted error is only standard error

Table A5. Half-life of  $^{236}\text{U}$

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^7$ y	Remarks
Jaffey et al. (1951) <sup>(40)</sup>	22.1-37.6	activity determ. by ion chamber energy analysis; U-determination by neutron induced fission counting	$2.46 \pm 0.06$	
Fleming et al. (1952) <sup>(27)</sup>	97.0	intermed. geom. $\alpha$ counting; gravimetric U-determination; chemical purity by spectrographic analysis	$2.391 \pm 0.018$	
Flynn et al. (1972) <sup>(41)</sup>	99.6	intermed. geom. $\alpha$ counting; U- determin. by titration; radiochem. purity by $\alpha$ -spectrometry	$2.3415 \pm 0.0014$	

Table A6. Half-life of  $^{238}\text{U}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^9$ y	Remarks
Kienberger (1949) <sup>(42)</sup> Kovarik and Adams (1955) <sup>(43)</sup>	depleted U nat U	$2\pi\alpha$ -counting; gravimetric U-determ. $\alpha$ -activity determin. by ion chamber; no specification about U determ.	$4.49 \pm 0.01$ $4.507 \pm 0.009$	
Leachman and Schmitt (1957) <sup>(44)</sup> <sup>(45)</sup> Steyn and Strelow (1960)	depleted U nat U	$2\pi\alpha$ -counting; U-determin. by gravi- metry and colometric analysis liq. scint. $\alpha$ counting (separation of $\beta$ -emitting daughters before counting and correction for growth); gravi- metric U-determin.	$4.56 \pm 0.03$	
Jaffey et al. (1971) <sup>(38)</sup>	> 99.9	intermed. geom. $\alpha$ -counting; U- determ. by titration; radiochem. purity by $\alpha$ -spectrom.	$4.4683 \pm 0.0024$	quoted error is only standard error

Table A7. Half-life of  $^{238}\text{Pu}$

Author (Year)	Material % isotope investig.	Method	Half-life in units of one y	Remarks
Hoffman et al. (1957) <sup>(46)</sup>		growth of $^{238}\text{Pu}$ from $^{242}\text{Cm}$ by $\alpha$ -counting	$86.4 \pm 0.3$	based on $T_{1/2}$ - $^{242}\text{Cm}$ 162.7d
Eichelberger et al. (1965) (47)		calorimetric determination	$87.60 \pm 0.06$	
Jordan (1967) (48)		calorimetric determination	$87.40 \pm 0.04$	should be cor- rected to ( $87.80 \pm 0.05$ ) (49)
Benson (1969) (50)		calorimetric determination	$87.75 \pm 0.05$	

Table A8. Half-life of  $^{239}\text{Pu}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^4$ y	Remarks
Stout and Jones (1947) <sup>(51)</sup>	> 99.9	calorimetric activity determ.; gravim. Pu-determ.	2.411 $\pm$ 0.024	
Cunningham and Werner (1949) <sup>(52)</sup>		2 $\pi\alpha$ -counting; gravimetric Pu-determ.	2.430 $\pm$ 0.037	
Westrum et al. (1949) <sup>(53)</sup>		2 $\pi\alpha$ -and low geom. $\alpha$ -counting; gravim. Pu-determ; chemical purity by spectrographic analysis	2.440 $\pm$ 0.070	
Wallmann (1951) <sup>(54)</sup>		low geom. $\alpha$ -counting; gravim. Pu-determ.; chem. purity by spectr. anal.; radioch. purity by $\alpha$ -spectrometry	2.436 $\pm$ 0.010	
Farwell et al. (1954) <sup>(55)</sup>		2 $\pi\alpha$ -counting; Pu-determ. by gravi- metry and titration	2.440 $\pm$ 0.048	
Markin (1959) <sup>(56)</sup>		low geom. $\alpha$ -counting; gravim. Pu- determ.	2.441 $\pm$ 0.003	
Dokuchaev (1959) <sup>(57)</sup>		low geom. $\alpha$ -counting; gravim. Pu- determ.; radiochem. purity by $\alpha$ -spectrometry	2.439 $\pm$ 0.003	
Ihle et al. (1966) <sup>(58)</sup>		liq. scint. $\alpha$ -counting; gravim. Pu determ.; radiochem. purity by $\alpha$ -spectrometry	2.435 $\pm$ 0.006	

continued on next page

Table A8-bis

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^4$ y	Remarks
Oetting (1968) <sup>(19)</sup>	94	calorimetric activity determ.; gravim. Pu determ.	$2.431 \pm 0.005$	corrected figure for the 1965- value of $(2.418 \pm 0.003) 10^4$ y (59)
Oetting (1971) <sup>(2)</sup>	> 99.9	calorimetric activity determ.; gravim. Pu determ. (electro refined Pu metal); chem. purity by spectro- graph. analysis; radiochem. purity by $\alpha$ -spectrometry	$2.4065 \pm 0.0050$	

Table A9. Half-life of  $^{240}\text{Pu}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^3$ y	Remarks
Westrum (1951) <sup>(60)</sup>	34.4	$2\pi\alpha$ -counting; gravim. Pu-determ.; chemical purity by spectrographic analysis	$6.24 \pm 0.12$	based on $T_{1/2}$ - $^{239}\text{Pu} = 2.44 \cdot 10^4$ y
Inghram et al. (1951) <sup>(61)</sup>		mass spectrometric determ. of $^{236}\text{U}/^{235}\text{U}$ (growing from $^{240}\text{Pu}$ and $^{239}\text{Pu}$ respectively)	$6.58 \pm 0.04$	
Wallmann (1951) <sup>(54)</sup>		low geom. $\alpha$ -counting; gravim. Pu determ.; spectrographic purity analysis; radiochem. purity by $\alpha$ -spectrometry	$6.76 \pm 0.03$	
Farwell et al. (1954) <sup>(55)</sup>		$2\pi\alpha$ -counting; Pu determ. by gravim. and titration	$6.30 \pm 0.60$	
Butler et al. (1956) <sup>(62)</sup>		$2\pi\alpha$ -counting (specific activity) $\alpha$ -energy analysis	$6.60 \pm 0.10$	
Dokuchaev (1959) <sup>(57)</sup>		low geom. $\alpha$ -counting; gravim. Pu- determ.; radiochemical analysis by $\alpha$ -spectrom.	$6.62 \pm 0.05$	
Oetting (1968) <sup>(19)</sup>	95.7	calorimetric activity determ.; gravim. Pu determ.	$6.524 \pm 0.010$	recalculated to $6.537 \cdot 10^3$ y by Schmorak <sup>(63)</sup> with $E(\alpha_o) =$ 5.1677 MeV

Table A10. Half-life of  $^{241}\text{Pu}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of one y	Remarks
Mac Kenzie et al. (1953) <sup>(64)</sup>		growth of $^{241}\text{Am}$ by $\alpha$ -spectrom.	14.1 $\pm$ 0.3	corrected by Ellis <sup>(9)</sup> on the basis $T_{1/2}^{241}\text{Am}$ = 433y
Rose and Milsted (1956) <sup>(65)</sup>	14.5	" " $\gamma$ -spectrom.	13.8 $\pm$ 0.3	
Brown et al. (1960) <sup>(66)</sup>	76.5	" " $\alpha$ -spectrom.	14.1 $\pm$ 0.3	
Smith (1961) <sup>(67)</sup>	96.2	" " "	13.9 $\pm$ 0.3	
Nisle and Stepan (1970) (68)	94.0	changes in reactivity	14.63 $\pm$ 0.27	
Shields (1970) <sup>(69)</sup>		changes in isotopic ratios by mass spectrometry	14.6 $\pm$ 0.4	cited by Ellis <sup>(9)</sup>
Cabell and Wilkins (1971) (70-71)	31.9	" " "	15.10 $\pm$ 0.14	1968-value was 14.98y <sup>(72)</sup>
Zeigler and Ferris (1973) (73)	3.0	" " "	14.89 $\pm$ 0.11	pooled value of observations in 6 laboratories



Table All. Half-life of  $^{242}\text{Pu}$

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^5$ y	Remarks
Butler et al. (1956) <sup>(76)</sup>		$^{242}\text{Pu}$ activity relative to $^{238}\text{Pu}$ by $\alpha$ -spectrometry	$3.65 \pm 0.05$	cited by Ellis <sup>(9)</sup> basis $T_{1/2} - ^{238}\text{Pu}$ = 87.75y
Butler et al. (1956) <sup>(62)</sup>	98.9	specific activity by ion chamber energy analysis	$3.79 \pm 0.05$	
Mech et al. (1956) <sup>(77)</sup>	96.3 - 98.7	$^{242}\text{Pu}$ activity relative to $^{240}\text{Pu}$ by $\alpha$ -spectrometry	$3.88 \pm 0.10$	basis $T_{1/2} - ^{240}\text{Pu}$ = $6.537 \cdot 10^3$ y
Bemis et al. (1969) <sup>(78)</sup>	99.9	$^{242}\text{Pu}$ activity relative to $^{239}\text{Pu}$ by $\alpha$ -spectrometry	$3.869 \pm 0.016$	basis $T_{1/2} - ^{239}\text{Pu}$ = $2.440 \cdot 10^4$ y
Durham and Malson (1970) <sup>(79)</sup>	90	$^{242}\text{Pu}$ activity relative to $^{238}\text{Pu}$ by $\alpha$ -spectrometry	$3.66 \pm 0.07$	basis $T_{1/2} - ^{238}\text{Pu}$ = 87.4y

Table A12. Half-life of  $^{244}\text{Pu}$

Author (Year)	Material % isotope investig.	Method	Half-life in units of $10^7$ y	Remarks
Diamond and Barnes (1956) (80)	0.05	separation of daughters; $\gamma$ -measure- ment	$7.6 \pm 2.0$	
Butler et al. (1956) (62)	0.08	separation of daughters; $\beta$ -measure- ment	$7.5 \pm 2.0$	
Fields et al. (1966) (81)	54 - 68	$^{244}\text{Pu}$ activity relative to $^{240}\text{Pu}$ and $^{242}\text{Pu}$ by $\alpha$ -spectrometry	$8.18 \pm 0.26$	basis $T_{1/2}^{240}\text{Pu}$ = $6.58 \cdot 10^3$ y and $T_{1/2}^{242}\text{Pu} =$ $3.79 \cdot 10^5$ y
Bemis et al. (1969) (78)	74.2-76.9	$^{244}\text{Pu}$ activity relative to $^{239}\text{Pu}$ by $\alpha$ -spectrometry	$8.28 \pm 0.10$	basis $T_{1/2}^{239}\text{Pu}$ = $2.440 \cdot 10^4$ y

Table A13. Half-life of  $^{241}\text{Am}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of one y	Remarks
Hall and Markin (1957) (3)		low geom. $\alpha$ counting; gravim. Am determ.; spectrographic purity analysis; radiochem. purity by $\alpha$ -spectrometry	$458.1 \pm 0.5$	
Wallmann et al. (1958) (4)		idem	$457.7 \pm 1.8$	
Oetting and Gun (1967) (6)		calorimetric activity determ.; gravim. Am determ.; chem. purity by spectrography and spark source mass spectrom.; radiochem. purity by $\alpha$ -spectrometry	$432.7 \pm 0.7$	
Stone and Hulet (1968) (7)		low geom. $\alpha$ -counting; gravim. Am determ.; spectrogr. purity analysis; radiochem. purity by $\alpha$ -spectrom.	$436.6 \pm 3.0$	same material as used by Oetting <sup>(6)</sup>
Brown and Probst (1968) (8)		low geom. $\alpha$ - and $2\pi\alpha$ -counting coulometric Am determ.	$433 \pm 7$	
Jove and Robert (1972) (10)		calorimetric activity determ.; gravim. Am determ.; spectro- graphic analysis; $\alpha$ -spectrometry	$426.3 \pm 2.1$	

Table A14. Half-life of  $^{252}\text{Cf}$ 

Author (Year)	Material % isotope investig.	Method	Half-life in units of y	Remarks
Eastwood et al. (1957) <sup>(82)</sup>	50	measurement of the decay of spontaneous fission (ion chamber) and $\alpha$ counting rate ( $2\pi\alpha$ )	$2.55 \pm 0.15$	
Metta et al. (1965) <sup>(83)</sup>	79.2	decay of spontaneous fission ( $2\pi$ -counting)	$2.646 \pm 0.004$	
De Volpi and Porges (1969) <sup>(84)</sup>		decay of neutron emission rate with a manganese bath facility	$2.621 \pm 0.006$	
Mijnheer and Van den Hauten (1973) <sup>(85)</sup>	50	idem	$2.659 \pm 0.010$	
Spiegel (1974)	79.3	idem	$2.638 \pm 0.007$	

# **NOTICE TO THE READER**

All scientific and technical reports published by the Commission of the European Communities are announced in the monthly periodical "euro-abstracts". For subscription (1 year: B.Fr. 1 025,—) or free specimen copies please write to :

**Office for Official Publications  
of the European Communities  
Boite postale 1003  
Luxembourg  
(Grand-Duchy of Luxembourg)**

# SALES OFFICES

The Office for Official Publications sells all documents published by the Commission of the European Communities at the addresses listed below, at the price given on cover. When ordering, specify clearly the exact reference and the title of the document.

## UNITED KINGDOM

*H.M. Stationery Office*  
P.O. Box 569  
London S.E. 1 — Tel. 01-928 69 77, ext. 365

## BELGIUM

*Moniteur belge — Belgisch Staatsblad*  
Rue de Louvain 40-42 — Leuvenseweg 40-42  
1000 Bruxelles — 1000 Brussel — Tel. 12 00 26  
CCP 50-80 — Postgiro 50-80

*Agency :*  
Librairie européenne — Europese Boekhandel  
Rue de la Loi 244 — Wetstraat 244  
1040 Bruxelles — 1040 Brussel

## DENMARK

*J.H. Schultz — Boghandel*  
Møntergade 19  
DK 1116 København K — Tel. 14 11 95

## FRANCE

*Service de vente en France des publications  
des Communautés européennes — Journal officiel*  
26, rue Desaix — 75 732 Paris - Cédex 15\*  
Tel. (1) 306 51 00 — CCP Paris 23-96

## GERMANY (FR)

*Verlag Bundesanzeiger*  
5 Köln 1 — Postfach 108 006  
Tel. (0221) 21 03 48  
Telex: Anzeiger Bonn 08 882 595  
Postscheckkonto 834 00 Köln

## GRAND DUCHY OF LUXEMBOURG

*Office for Official Publications  
of the European Communities*  
Boîte postale 1003 — Luxembourg  
Tel. 4 79 41 — CCP 191-90  
Compte courant bancaire: BIL 8-109/6003/200

## IRELAND

*Stationery Office — The Controller*  
Beggars Bush  
Dublin 4 — Tel. 6 54 01

## ITALY

*Libreria dello Stato*  
Piazza G. Verdi 10  
00198 Roma — Tel. (6) 85 08  
CCP 1/2640

## NETHERLANDS

*Staatsdrukkerij- en uitgeverijbedrijf*  
Christoffel Plantijnstraat  
's-Gravenhage — Tel. (070) 81 45 11  
Postgiro 42 53 00

## UNITED STATES OF AMERICA

*European Community Information Service*  
2100 M Street, N.W.  
Suite 707  
Washington, D.C., 20 037 — Tel. 296 51 31

## SWITZERLAND

*Librairie Payot*  
6, rue Grenus  
1211 Genève — Tel. 31 89 50  
CCP 12-236 Genève

## SWEDEN

*Librairie C.E. Fritze*  
2, Fredsgatan  
Stockholm 16  
Post Giro 193, Bank Giro 73/4015

## SPAIN

*Libreria Mundi-Prensa*  
Castello 37  
Madrid 1 — Tel. 275 51 31

## OTHER COUNTRIES

*Office for Official Publications  
of the European Communities*  
Boîte postale 1003 — Luxembourg  
Tel. 4 79 41 — CCP 191-90  
Compte courant bancaire: BIL 8-109/6003/200